

# Rules of Numerical Integration in Details

VISHAL VAMAN MEHTRE<sup>1</sup>, KETANSHU KAILASH SHAKYA<sup>2</sup>

<sup>1</sup> Assistant Professor, Department of Electrical Engineering, Bharati Vidyapeeth, University College of Engineering, PUNE

<sup>2</sup> Department of Electrical Engineering, Bharati Vidyapeeth, University College of Engineering, PUNE

**Abstract --** In this review paper we have tried to write all the points related to the numerical integration. Numerical integration are not the normals integration formulas but they are solved by writing different formulas which are not the general integration formulas. They are simply the algebraic formulas which can be solved easily, although they are quite leandy. There are three ways to solve the definite integrals (in which limits are given) -

1. Trapezoidal Rule
2. Simpson's 1/3rd Rule
3. Simpson's 3/8th Rule

In this paper we have tried our best for the explanation of all these three topics and also have given example for further understanding.

## I. INTRODUCTION

Integration is the method of finding the area of the function plot on the graph.

**Trapezoidal rule** - A trapezoidal rule is done by the approximation of the region which is covered by the graph of the function  $f(x)$  as a trapezoid and further measuring its area. In this the interval is broken into subintervals and then chained as a sum to get the appropriate definite integral answer. It is based on the approximation of integral by a first order polynomial [6].

Its equation is given as -

$$I = h/2 [(y_0 + y_n) + 2(y_1 + y_2 + y_3 + \dots + y_{n-1})]$$

**Simpson's 1/3<sup>rd</sup> rule** – It is similar to the trapezoidal rule which is used to calculate the area of a given function by the definite integral form. Only the difference is that it is the extended form of trapezoidal rule as it is used to calculate the second order polynomial [1].

Its equation is given as -

$$I = h/3 [(y_0 + y_n) + 4(y_1 + y_3 + y_5 + \dots) + 2(y_2 + y_4 + y_6 + \dots)]$$

**Simpson's 3/8<sup>th</sup> rule** - Simpson's 3/8 rule is designed to calculate the third order (cubic) polynomial function.

Its equation is given as -

$$I = 3h/8[(y_0 + y_n) + 3(y_1 + y_2 + y_4 + y_5 + y_7 + \dots) + 2(y_3 + y_6 + y_9 + \dots)]$$

**Trapezoidal Rule** - Let the function be given  $f(x)$  whose integration we have to calculate from  $x_0$  to  $x_n$ . The function is given as  $y = f(x)$ .

Now, the function according to the limits are given as -

$$Y_0 = f(x_0) \text{ and}$$

$$Y_n = f(x_n).$$

$$f(x) = f(x_0) + f(x_n)$$

$$I \text{ (whose limit is from } x_0 \text{ to } x_n) = \int [ f(x_0) + \{ (f(x_n) - f(x_0)) (x - x_0) \} / (x_n - x_0) ] dx$$

$$I \text{ (whose limit is from } x_0 \text{ to } x_n) = \int [ f(x_0) + \{ xf(x_n) - xf(x_0) \} / (x_n - x_0) - \{ f(x_n)x_0 - f(x_0)x_n \} / (x_n - x_0) ] dx$$

$$I \text{ (whose limit is from } x_0 \text{ to } x_n) = \int [ \{ x_n f(x_0) - x_0 f(x_0) + xf(x_n) - xf(x_0) - x_0 f(x_n) + x_0 f(x_0) \} / (x_n - x_0) ] dx$$

$$I \text{ (whose limit is from } x_0 \text{ to } x_n) = \int [ \{ x_n f(x_0) + xf(x_n) - xf(x_0) - x_0 f(x_n) \} / (x_n - x_0) ] dx$$

$$I \text{ (putting both upper and lower limits)} = x_n f(x_0) - x_0 f(x_n) + \{ (x_n + x_0)(f(x_n) - f(x_0)) \} / 2$$

$$I = \{ 2x_n f(x_0) - 2x_0 f(x_n) + x_n f(x_n) - x_n f(x_0) + x_0 f(x_n) - x_0 f(x_0) \} / 2$$

$$I = \{ x_n f(x_0) - x_0 f(x_n) + x_n f(x_n) - x_0 f(x_0) \} / 2$$

$$I = \{ x_n ( f(x_0) + f(x_n) ) - x_0 ( f(x_n) + f(x_0) ) \} / 2$$

$$I = ( x_n - x_0 ) ( f(x_n) + f(x_0) ) / 2$$

$$I = ( x_n - x_0 ) ( y_n + y_0 ) / 2$$

I = width \* average of y coordinate

**Composite Integral –**

$$\int(x_0 \text{ to } x_n) y \, dx = \int(x_0 \text{ to } x_1) y \, dx + \int(x_1 \text{ to } x_2) y \, dx + \int(x_2 \text{ to } x_3) y \, dx + \dots$$

$$I = [ (x_1 - x_0)(y_1 + y_0)/2 + (x_2 - x_1)(y_2 + y_1)/2 + (x_3 - x_2)(y_3 + y_2)/2 + \dots ]$$

{ Since  $(x_1 - x_0) = (x_2 - x_1) = (x_3 - x_2) = h$  }

$$I = \int(x_0 \text{ to } x_n) y \, dx = h [(y_1 + y_0)/2 + (y_2 + y_1)/2 + (y_3 + y_2)/2 + \dots ]$$

$$I = h [ y_0/2 + y_1 + y_2 + y_3 + \dots + y_n/2 ]$$

$$I = h/2 [ (y_0 + y_n) + 2(y_1 + y_2 + y_3 + \dots + y_{n-1}) ]$$

Is the required trapezoidal rule.

**SIMPSON'S 1/3<sup>rd</sup> RULE –**

It is the 3 points and 2<sup>nd</sup> degree equation.

It has three points A(x<sub>0</sub>, y<sub>0</sub>), B(x<sub>k</sub>, y<sub>k</sub>), C(x<sub>n</sub>, y<sub>n</sub>)

Let the equation of parabola is given as y = f(x)[5].

Now,

By lagrange's interpolation method-

$$Y = f(x) = \{ (x - x_k)(x - x_n)/(x_0 - x_k)(x_0 - x_n) \} * y_0 + \{ (x - x_0)(x - x_n) / (x_k - x_0)(x_k - x_n) \} * y_k + \{ (x - x_0)(x - x_k)/(x_n - x_0)(x_n - x_k) \} * y_n$$

$$\int(x_0 \text{ to } x_n) y \, dx = \int(x_0 \text{ to } x_n) [ \{ (x^2 - xx_n - xx_k + x_n x_k) / (x_0 - x_k)(x_0 - x_n) \} * y_0 + \{ (x^2 - xx_n - xx_0 + x_0 x_n) / (x_k - x_0)(x_k - x_n) \} * y_k + \{ (x^2 - xx_k - xx_0 - x_0 x_k) / (x_n - x_0)(x_n - x_k) \} * y_n ] \, dx$$

$$\int(x_0 \text{ to } x_n) y \, dx = \int(x_0 \text{ to } x_n) [ \{ (x^2 - (x_n + x_k)x + x_n x_k) / (x_0 - x_k)(x_0 - x_n) \} * y_0 + \{ (x^2 - (x_n + x_0)x + x_0 x_k) / (x_k - x_0)(x_k - x_n) \} * y_k + \{ (x^2 - (x_n + x_0)x + x_0 x_k) / (x_n - x_0)(x_n - x_k) \} * y_n ] \, dx$$

$$x_n)/(x_k - x_0)(x_k - x_n) \} * y_k + \{ (x^2 - (x_k + x_0)x + x_0 x_k) / (x_n - x_0)(x_n - x_k) \} * y_n ] \, dx$$

$$I = y_0 / (x_0 - x_k)(x_0 - x_n) [ (x_n^3 - x_0^3) / 3 - (x_n + x_k)(x_n^2 - x_0^2) / 2 + x_n x_k (x_n - x_0) ] + y_k / (x_k - x_0)(x_k - x_n) [ (x_n^3 - x_0^3) / 3 - (x_n + x_0)(x_n^2 - x_0^2) / 2 + x_0 x_n (x_n - x_0) ] + y_n / (x_n - x_0)(x_n - x_k) [ (x_n^3 - x_0^3) / 3 - (x_k + x_0)(x_n^2 - x_0^2) / 2 + x_0 x_k (x_n - x_0) ]$$

$$\int(x_0 \text{ to } x_n) y \, dx = (x_n - x_0) / 6 [ y_0 + 4y_k + 2y_n ] = h/3 (y_0 + 4y_k + 2y_n) = h/3 (y_0 + 4y_1 + 2y_2) + h/3 (y_2 + 4y_3 + 2y_4) + h/3 (y_4 + 4y_5 + 2y_6)$$

$$I = h/3 [ (y_0 + y_n) + 4(y_1 + y_3 + y_5 + \dots) + 2(y_2 + y_4 + y_6 + \dots) ]$$

Therefore, this is the required simpson's 1/3<sup>rd</sup> rule.

**SIMPSON'S 3/8<sup>th</sup> RULE –**

Taking four points (x<sub>0</sub>,y<sub>0</sub>), (x<sub>1</sub>,y<sub>1</sub>), (x<sub>2</sub>,y<sub>2</sub>) and (x<sub>3</sub>,y<sub>3</sub>) Now, finding the integration of function y from x<sub>0</sub> to x<sub>3</sub> we get-

$$\int(x_0 \text{ to } x_3) y \, dx = [ \{ (x - x_1)(x - x_2)(x - x_3) / (x_0 - x_1)(x_0 - x_2)(x_0 - x_3) \} * y_0 + \{ (x - x_0)(x - x_2)(x - x_3) / (x_1 - x_0)(x_1 - x_2)(x_1 - x_3) \} * y_1 + \{ (x - x_0)(x - x_1)(x - x_3) / (x_2 - x_0)(x_2 - x_1)(x_2 - x_3) \} * y_2 + \{ (x - x_0)(x - x_1)(x - x_2) / (x_3 - x_0)(x_3 - x_1)(x_3 - x_2) \} * y_3 ] \, dx$$

On solving this equation we get-

$$I = (x_3 - x_0) / 8 (y_0 + 3y_1 + 3y_2 + 2y_3)$$

Here we know that h = x<sub>1</sub> - x<sub>0</sub> = x<sub>2</sub> - x<sub>1</sub> = x<sub>3</sub> - x<sub>2</sub>

Therefore,

$$\int(x_0 \text{ to } x_n) y \, dx = 3h/8 * [ (y_0 + y_n) + 3(y_1 + y_2 + y_4 + y_5 + y_7 + \dots) + 2(y_3 + y_6 + y_9 + \dots) ]$$

Is the required simpson's 3/8<sup>th</sup> rule.

**EXAMPLES:**

Que. 01. Evaluate integration  $\int(0 \text{ to } 0.9) \log_e(1 + \sqrt{x}) \, dx$  using trapezoidal rule with 9 subinterval.

Ans. Here x<sub>0</sub> = 0, x<sub>n</sub> = 0.9 and n = 9

Now, h = (0.9 - 0) / 9

$$= 0.1$$

x	0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
y	0	0.274	0.369	0.436	0.490	0.534	0.573	0.607	0.638	0.667

Now by trapezoidal rule-

$$I = 0.1/2 [(0 + 0.667) + 2(0.274 + 0.369 + 0.436 + 0.490 + 0.534 + 0.573 + 0.607 + 0.638) ]$$

Therefore I = 0.42545 is the integration.

Que. 02. Evaluate  $\int(1.0 \text{ to } 1.8) (e^x + e^{-x})/2 \text{ dx}$  using simpson's 1/3<sup>rd</sup> rule

By taking h = 0.2.

Ans. Here  $x_0 = 1.0$ ,  $x_n = 1.8$  and  $h = 0.2$

$$\text{Now, } n = (1.8 - 1.0)/ 0.2 = 4$$

X	1.0	1.2	1.4	1.6	1.8
y	1.5430	1.8106	2.1508	2.5774	3.1074

Now by simpson's 1/3<sup>rd</sup> rule -

$$I = 0.2/3 [ (1.5430 + 3.1074) + 4(1.8106 + 2.5774) + 2(2.1508) ]$$

Therefore I = 1.764 is the integration.

Que. 03. Evaluate  $\int(0 \text{ to } \pi/2) e^{\sin x} \text{ dx}$  using simpson's 3/8<sup>th</sup> rule with 6 subintervals.

Ans. Here,  $x_0 = 0$ ,  $x_n = \pi/2$  and  $n = 6$

$$\text{Now, } h = (\pi/2 - 0) / 6 = \pi/12$$

X	0	$\pi/12$	$\pi/6$	$\pi/4$	$\pi/3$	$5\pi/12$	$\pi/2$
Y	1	1.2951	1.6483	2.0275	2.3768	2.6267	2.7182

Now by simpson's 3/8<sup>th</sup> rule-

$$I = 3\pi/12*8 [(1 + 2.7182) + 3(1.2951 + 1.6483 + 2.3768 + 2.6267) + 2 * 2.0275]$$

Therefore I = 3.1021 is the integration.

### II. APPLICATIONS

- By these method we can find extremely accurate answer when the function is periodic i.e. the interval is periodic.

- By this method there is faster convergence of the function (i.e. they can be calculated faster without the remembrance of the formula).

### III. CONCLUSION

By this paper we come to know about the different ways which are helpful to calculate the numerical integrals, their formulas and the applications. Examples are also helpful to understand the topic.

### ACKNOWLEDGEMENT

We would like to express our special thanks of gratitude to **Dr. D. S Bankar** Head of Department of Electrical Engineering for their able guidance and support for completing my research paper. I would also like to thank the faculty members of the Department of Electrical Engineering would helped us with extended support.

### REFERENCES

- [1] S. S. Sastry, "Introductory Methods of Numerical Analysis", 4<sup>th</sup> Edition, PHI
- [2] Dr. J. S. Chitodia, "Numerical Methods", Technical Publications
- [3] M. K. Jain, R. K. Jain, S. R. K. Iyengar, "Numerical Methods for Scientific and Engineering Computation", 6<sup>th</sup> Edition, New Age International Publishers
- [4] Felix O. Mettle, "A proposed method for Numerical Integration", 'British Journal of Mathematics and computer science', 31<sup>th</sup> may 2016.
- [5] Louis A. Talman."Simpson's Rule is Exact For Quintics", 'The American Mathematical Monthly', Feb 2006
- [6] Lloyd N. Trefethen," The Exponentially Convergent Trapezoidal Rule", 'SIAM', 2014